

Aerospace Europe Conference 2023

Joint 10th EUCASS – 9th CEAS Conference

Abstract #XXX (to be filled by the organizers)
Preferred Topics: CFDMPS / AEROFLIPHY
Corresponding author: Aaron Nagel
e-mail of corresponding author: aaron.nagel@dlr.de
Type: Oral
Status of corresponding author: Student

Title

Modeling of polyatomic gas flows within a kinetic Fokker-Planck approach using a direct modeling method

Authors

Aaron NAGEL ^{1*}, Leo Basov ², Martin Grabe ³

* Corresponding author

¹ German Aerospace Center (DLR), Bunsenstraße 10, D-37073 Göttingen, Germany, aaron.nagel@dlr.de

² German Aerospace Center (DLR), Bunsenstraße 10, D-37073 Göttingen, Germany, leo.basov@dlr.de

³ German Aerospace Center (DLR), Bunsenstraße 10, D-37073 Göttingen, Germany, martin.grabe@dlr.de

Abstract

The modeling of gas flows as for re-entry vehicles or jet plume expansion faces a large range of the Knudsen number. With Bird's DSMC method [1], the Boltzmann equation, describing the evolution of the system by a probability density function, can be solved accurately. However, the computational expenses become unfeasible for small Knudsen numbers due to the complexity of the Boltzmann collision operator with the large amount of collisions. Therefore, the complex collision operator in the Boltzmann equation is approximated by a Fokker-Planck (FP) operator in velocity space. While DSMC models pairwise collisions, the collisions by the FP operator are modeled by local drift and diffusion coefficients, that are matched to reproduce the production terms of the Boltzmann collision operator in the continuum limit [2]. Both methods use computational particles. This way, a hybrid modeling approach can be set up where FP can be used in regions of high densities and small Knudsen numbers, while DSMC is used for rarefied flow regions and large Knudsen numbers [3].

In contrast to monatomic gas flows, polyatomic molecules can take up a significant amount of energy in internal modes like rotational and vibrational, which have a large influence on the entire flow field. An extension of the FP operator to a diatomic modeling has been approached using two methods: the Master-equation ansatz [4] and a direct modeling approach [5]. In the meantime, the Master-equation ansatz has been extended to a polyatomic model [6], while a polyatomic extension of the direct modeling approach will be presented in this paper.

Comparisons with the Master-equation ansatz in terms of the quality of the flow field and flow properties as well as the overall performance are made. Finally, temporal relaxations of energies are investigated.

References

- [1] G. A. Bird, Molecular Gas Dynamics and the direct Simulation of Gas Flows (Oxford University Press, New York, 1994).
- [2] M. H. Gorji, M. Torrilhon, P. Jenny, Journal of fluid mechanics 680, pp. 574–601 (2011).
- [3] M. H. Gorji, P. Jenny, Journal of Computational Physics 287, pp. 110–129 (2015).
- [4] C. Hepp, M. Grabe, K. Hannemann, Journal of Computational Physics 418, 109638 (2020).
- [5] M. H. Gorji, P. Jenny, Physics of fluids 25.6, p. 062002 (2013).
- [6] L. Basov, M. Grabe, 32nd International Symposium on Rarefied Gas Dynamics (2022) - (under review).