

Aerospace Europe Conference 2023

Joint 10th EUCASS – 9th CEAS Conference

Abstract #XXX (to be filled by the organizers)

Preferred Topics: CFDMPS (3 maximum from the list of topics)

Corresponding author: Michele Capriati

e-mail of corresponding author: michele.capriati@vki.ac.be

Type: Oral

Status of corresponding author: Student

For student corresponding author: student member of one of the following: no

Title

Bayesian calibration of a finite-rate nitridation model from molecular beam and plasma wind tunnel experiments

Authors

Michele Capriati^{a,b*}, Anabel del Val^b, Thomas Schwartzentruber^c, Timothy Minton^d, Pietro Congedo^a, Thierry Magin^{b,e}

^a Inria, Centre de Mathématiques Appliquées, Ecole Polytechnique, IPP, Route de Saclay, 91120 Palaiseau, France

^b von Karman Institute for Fluid Dynamics, Chaussée de Waterloo 72, 1640 Rhode-Saint-Genese, Belgium

^c University of Minnesota, 110 Union Street SE, 55455 Minneapolis, Minnesota (USA)

^d University of Colorado, 3775 Discovery Dr, 80303, Boulder, Colorado (USA)

^e Aero-Thermo-Mechanics Laboratory, École Polytechnique de Bruxelles, Université Libre de Bruxelles, 91050 Brussels, Belgium

Abstract

The accurate modeling of gas-surface interaction phenomena is crucial for the prediction of the heat flux and the mass loss experienced by atmospheric entry bodies flying at hypersonic speeds. Gas-surface interactions refer to the set of phenomena occurring between the reactive gas and the material surface, involving heterogeneous chemical reactions together with mass and heat exchange. The surface chemical production rates can be computed using phenomenological or finite-rate chemistry models. In [1] the finite-rate chemistry model derived by Prata et al. [2] was employed to numerically simulate a low-pressure nitridation experimental campaign carried out at the von Karman Institute Plasmatron facility. The model was calibrated both on molecular beam data and on Plasmatron data. The work did not account for experimental as well as parametric uncertainties. In turn, some observed features (e.g., surface recession rates) were not in agreement with the experimental values. This issue underlined the necessity to account for experimental as well as parametric uncertainties during the calibration of the model parameters. In [3], it was shown that Bayesian analyses, which leverage experimental and parametric uncertainties, offer a suitable framework for the calibration of finite-rate surface chemistry models. In this work, we propose to calibrate nitridation surface chemical rates in a Bayesian framework. The calibration is performed by combining wind tunnel and molecular beam data, a first in the ablation literature. The plasma wind tunnel data are explained using a surrogate model that captures the underlying boundary layer in front of the test sample. Such a surrogate model is trained on CFD computations to accurately describe recession rates and CN densities on the stagnation line. The molecular beam data are explained through an analytical model, as in [2]. The parameters' posterior distribution is obtained by constructing a Markov Chain Monte Carlo. Such calibration allows the model to be consistent with both molecular beam and plasma wind tunnel data.

References

- [1] Capriati, M. and Prata, K. and Schwartzentruber, T. and Candler, G. and Magin, T., Development of a nitridation gas-surface boundary condition for high-fidelity hypersonic simulations. 14th WCCM-ECCOMAS Congress, CIMNE, 2021.
- [2] Prata, K. S. and Schwartzentruber, T. E. and Minton, T. K. Air-Carbon Ablation Model for Hypersonic Flight from Molecular Beam Data. AIAA Journal, 2022.
- [3] del Val, A, Le Maître, O., Chazot, O. and Congedo, P.M. and Magin, T. Stochastic calibration of a carbon nitridation model from plasma wind tunnel experiments using a Bayesian formulation. Carbon, 2022.